

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 389 282
A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90303093.0

(51) Int. Cl.⁵: C07D 473/06, A61K 31/52

(22) Date of filing: 22.03.90

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

(30) Priority: 23.03.89 GB 8906792

(43) Date of publication of application:
26.09.90 Bulletin 90/39

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: BEECHAM - WUELFING GmbH & Co.
KG
Stresemannallee 6 P.O. Box 25
D-4040 Neuss(DE)

Applicant: Beecham Group p.l.c.
SB House Great West Road
Brentford Middlesex TW8 9BD(GB)

(72) Inventor: Maschler, Harald, c/o
Beecham-Wuelfing
GmbH & Co KG, Bethelner Landstrasse
D-3212 Gonau (Leine)(DE)

Inventor: Spicer, Barbara Ann, Beecham
Pharmaceuticals

Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT 18 5XQ(GB)


Inventor: Smith, Harry, Beecham
Pharmaceuticals

Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT 18 5XQ(GB)

(74) Representative: Rutter, Keith et al
Smith Kline Beecham, Corporate Patents,
Great Burgh, Yew Tree Bottom Road
Epsom Surrey KT18 5XQ(GB)

(54) Xanthinederivatives, process for their preparation and their pharmaceutical use.

(57) A method for the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or other disorders which method comprises the administration of an effective, non-toxic amount of a compound of formula (I):



(I)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R¹ and R² each independently represent alkyl or a moiety of formula (a):

-(CH₂)_m-A (a)

wherein m represents zero or an integer 1, 2 or 3;

A represents a substituted or unsubstituted cyclic hydrocarbon radical; and

R³ represents a halogen atom, a nitro group, or a group -NR⁴R⁵ wherein R⁴ and R⁵ each independently represents hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are

EP 0 389 282 A2

EP 0 389 282 A2

attached forming an optionally substituted, heterocyclic group; certain novel compounds falling within formula (I) and compositions comprising such compounds.

TREATMENT AND COMPOUNDS

The present invention relates to a novel method of treatment and to certain novel compounds having pharmacological activity, to a process for the preparation of such compounds, to pharmaceutical compositions containing such compounds and to the use of such compounds and compositions in medicine.

Molecular Pharmacology, Volume 6, No. 6, 1970, p.597-603 discloses 1,3-dimethyl-8-nitro-xanthine.
 5 This compound is disclosed as having lipolytic activity.

Annalen der Chemie, 47, 362-365 (1957) discloses 1,3-dimethyl-8-amino-xanthine and a process by which it may be prepared. No pharmacological utility is disclosed for this compound.

Drug Res. 27(1) Nr 19, 1977, pages 4-14, Van K.H. Klingler discloses certain 1,3-dimethyl- 8-substituted xanthines as intermediates solely in the synthesis of phenylethyl aminoalkyl xanthines.

10 Drug Res. 31 (11), Nr. 12, 1981, R.G. Werner et al, pages 2044-2048 discloses certain 1,3-dimethyl-8-substituted xanthines. No pharmacological activity is disclosed for these compounds.

It has now been discovered that certain 8-substituted xanthines have a protective effect against the consequences of cerebral metabolic inhibition. The said compounds improve data acquisition or retrieval following transient forebrain ischaemia and are therefore useful in the treatment of cerebral vascular and
 15 neuronal degenerative disorders associated with learning, memory and cognitive dysfunctions including cerebral senility, multi-infarct dementia, senile dementia of the Alzheimer type, age associated memory impairment and certain disorders associated with Parkinson's disease.

These compounds are also indicated to have neuroprotectant activity. They are therefore useful in the prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events, including
 20 cerebral ischaemia due to cardiac arrest, stroke and also after cerebral ischaemic events such as those resulting from surgery and/or during childbirth. In addition treatment with the compound is indicated to be of benefit for the treatment of functional disorders resulting from disturbed brain function following ischaemia.

These compounds are also active in increasing the oxygen tension in ischaemic skeletal muscle. This property results in an increase in the nutritional blood flow through ischaemic skeletal muscle which in turn
 25 indicates that the compounds of the invention are of potential use as agents for the treatment of peripheral vascular disease such as intermittent claudication.

These compounds also act as phosphodiesterase inhibitors and elevate cyclic AMP levels and are therefore of potential use in the treatment of proliferative skin disease in human or non-human mammals.


These compounds are also indicated to have bronchodilator activity and thus to be of potential use in
 30 the treatment of disorders of the respiratory tract, such as reversible airways obstruction and asthma.

It has now also surprisingly been discovered that these compounds are good inhibitors of induced blood eosinophilia and that they are therefore potentially useful in the treatment and/or prophylaxis of disorders associated with increased numbers of eosinophils, such as asthma, and allergic disorders associated with atopy, such as urticaria, eczema and rhinitis.

35 Certain of the novel compounds are also indicated to possess useful adenosine A1 antagonist activity.

Finally the present compounds also show good metabolic stability.

Accordingly, the invention provides a method for the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events and/or peripheral vascular disease and/or proliferative skin disease
 40 and/or for disorders of the respiratory tract and/or the treatment or prophylaxis of disorders associated with increased numbers of eosinophils and allergic disorders associated with atopy, which method comprises the administration of an effective, non-toxic amount of a compound of formula (I):



(I)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R¹ and R² each independently represent alkyl or a moiety of formula (a):

-(CH₂)_m-A (a)

wherein m represents zero or an integer 1, 2 or 3;

A represents a substituted or unsubstituted cyclic hydrocarbon radical; and

R³ represents a halogen atom, a nitro group, or a group -NR⁴R⁵ wherein R⁴ and R⁵ each independently represents hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are attached form an optionally substituted, heterocyclic group; to a human or non-human mammal in need thereof.

In another aspect, the invention provides the use of a compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events and/or peripheral vascular disease and/or proliferative skin diseases and/or disorders of the respiratory tract and/or the treatment or prophylaxis of disorders associated with increased numbers of eosinophils and allergic disorders associated with atopy.

The present invention also provides a pharmaceutical composition comprising a compound of formula (I) or if appropriate a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier therefor.


The present invention also provides a pharmaceutical composition comprising a compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, providing that in the compound of formula (I) when R¹ and R² both represent methyl then R³ is not a nitro group, and a pharmaceutically acceptable carrier therefor.

In a further aspect the invention provides a compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, providing that in the compound of formula (I) when R¹ and R² both represent methyl then R³ is not a nitro group, for use as an active therapeutic substance.

The invention also provides a compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, providing that in the compound of formula (I) when R¹ and R² both represent methyl then R³ is not a nitro group, for use in the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events and/or peripheral vascular disease and/or proliferative skin diseases and/or disorders of the respiratory tract and/or the treatment or prophylaxis of disorders associated with increased numbers of eosinophils and allergic disorders associated with atopy.

As indicated above, certain of the compounds of formula (I) are novel and form a further aspect of the present invention.

Accordingly, the invention also provides a compound of formula (IA):



(IA)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R¹ and R² each independently represent alkyl or a moiety of formula (a):

-(CH₂)_m-A (a)

wherein m represents zero or an integer 1, 2 or 3, A represents a substituted or unsubstituted cyclic hydrocarbon radical, providing that when R¹ represents methyl then R² is not methyl; and

R^{3a} represents a halogen atom, a nitro group, or a group -NR⁴R⁵ wherein R⁴ and R⁵ each independently represent hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are attached form an optionally substituted heterocyclic group.

Suitably, R¹ represents a moiety of formula (a).

Suitably, R² represents a moiety of formula (a).

Preferably, R¹ and R² each independently represent a moiety of formula (a).

Suitably, A is unsubstituted. Favourably, A represents a substituted or unsubstituted C₃₋₈ cycloalkyl group, especially a C₃₋₆ cycloalkyl group.

In particular, A represents a substituted or, preferably, unsubstituted cyclopropyl, cyclobutyl, cyclopentyl,

tyl or cyclohexyl group.

Favourably, A represents a cyclopropyl group or a cyclobutyl group.

Preferably, A represents a cyclopropyl group.

When R¹ or R² represents alkyl, a preferred alkyl group is an n-butyl group.

5 An example of R³ or R^{3a} includes a nitro group or a group -NHR⁴ wherein R⁴ represents hydrogen or alkylcarbonyl.

When R³ or R^{3a} represents a halogen atom it is suitably a bromine or a chlorine atom.

When either of R⁴ or R⁵ represents alkyl or alkylcarbonyl, it is preferred if the other of R⁴ or R⁵ represents hydrogen.

10 An example of an alkylcarbonyl group is an acetyl group.

Suitable heterocyclic groups include saturated or unsaturated heterocyclic groups having single or fused rings, each ring having 5 to 7 ring atoms which ring atoms optionally comprise up to two additional hetero atoms selected from O, N or S.

Favoured heterocyclic groups include rings comprising 5 to 7, especially 5 or 6 and preferably 6, ring
15 atoms.

Favoured additional hetero atoms are O or N, especially O.

Favoured heterocyclic groups are saturated heterocyclic groups.

Favoured heterocyclic groups are single ring heterocyclic groups.

Favoured heterocyclic groups comprising 5 ring atoms include pyrrolidinyl groups.

20 Favoured heterocyclic groups comprising 6 ring atoms include piperidinyl or morpholinyl groups.

Suitably, R³ represents amino.

Suitably, R^{3a} represents amino.

Suitably, m represents zero or the integer 1.

Favourably, m represents 1.

25 Suitable pharmaceutically acceptable salts are pharmaceutically acceptable base salts and pharmaceutically acceptable acid addition salts. Generally compounds of formula (I) wherein R³ is nitro form base salts, suitable pharmaceutically acceptable base salts of the compounds of formula (I) include 7-N base salts including metal salts, such as alkali metal salts for example sodium salts, or organic amine salts such as that provided with ethylenediamine.

30 Certain of the compounds of formula (I) wherein R³ is amino form acid addition salts, suitable acid addition salts of the compounds of formula (I) are the acid addition salts including pharmaceutically acceptable inorganic salts such as the sulphate, nitrate, phosphate, borate, hydrochloride and hydrobromide and pharmaceutically acceptable organic acid addition salts such as acetate, tartrate, maleate, citrate, succinate, benzoate, ascorbate, methane-sulphate, α -keto glutarate, α -glycerophosphate and glucose-1-
35 phosphate. Preferably the acid addition salt is a hydrochloride salt.

The pharmaceutically acceptable salts of the compounds of formula (I) are prepared using conventional procedures.

A suitable compound of formula (I) is a compound of formula (IA).

40 When used herein the term 'cyclic hydrocarbon radical' includes single ring and fused ring, cyclic hydrocarbons comprising up to 8 carbon atoms in each ring, suitably up to 6 carbon atoms, for example 3, 4, 5 or 6 carbon atoms.


Suitable optional substituents for any cyclic hydrocarbon radical includes a C₁₋₆ alkyl group or a halogen atom.

45 When used herein the term 'alkyl' whether used alone or when used as part of another group (for example as in an alkylcarbonyl group) includes straight and branched chain alkyl groups, containing from 1 to 12 carbon atoms, suitably 1 to 6 carbon atoms, for example methyl, ethyl, propyl or butyl.

50 When used herein the expression 'proliferative skin diseases' means benign and malignant proliferative skin diseases which are characterized by accelerated cell division in the epidermis, dermis or appendages thereto, associated with incomplete tissue differentiation. Such diseases include: psoriasis, atopic dermatitis, non-specific dermatitis, primary irritant contact dermatitis, allergic contact dermatitis, basal and squamous cell carcinomas of the skin, lamellar ichthyosis, epidermolytic hyperkeratosis, premalignant sun induced keratosis, non-malignant keratosis, acne, and seborrheic dermatitis in humans and atopic dermatitis and mange in domesticated animals.

55 The compounds of formula (I) are preferably in pharmaceutically acceptable form. By pharmaceutically acceptable form is meant, inter alia, of a pharmaceutically acceptable level of purity excluding normal pharmaceutical additives such as diluents and carriers, and including no material considered toxic at normal dosage levels. A pharmaceutically acceptable level of purity will generally be at least 50% excluding normal pharmaceutical additives, preferably 75%, more preferably 90% and still more preferably 95%.

The invention further provides a process for the preparation of a compound of formula (IA), which process comprises reacting a compound of formula (II):



(II)

wherein R^{1a} represents R^1 , as defined in relation to formula (IA), or a group convertible to R^1 and R^{2a} represents R^2 , as defined in relation to formula (IA), or a group convertible thereto, with a reagent capable of substituting the C-8 hydrogen of the compound of formula (II) with a group R^{3b} wherein R^{3b} represents R^{3a} , as defined above in relation to formula (IA), or a group convertible thereto; and thereafter, if required carrying out one or more of the following optional steps:

- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) when R^{3b} is not R^{3a} , converting R^{3b} to R^{3a} ;
- (iii) converting a compound of formula (IA) into a further compound of formula (IA);
- (iv) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

For compounds of formula (IA) wherein R^{3a} represents nitro, R^{3b} preferably represents R^{3a} i.e. nitro.

For compounds of formula (IA) wherein R^{3a} represents other than nitro, R^{3b} preferably represents a group convertible to R^{3a} .

One preferred group R^{3b} is a nitro group which may then if required be converted to a group R^{3a} .

Suitable reagents for substituting the C-8 hydrogen of the compound of formula (II) with a group R^{3b} are the appropriate conventional reagents.

The conditions of reaction for the substitution of the C-8 hydrogen of the compound of formula (II) will of course depend upon the particular reagent chosen, and in general the conditions used will be those which are conventional for the reagent used.


One particularly suitable reagent is a nitrating agent.

In one convenient form of the abovementioned process the compound of formula (II) is reacted with a suitable nitrating agent to provide a compound of formula (IA) wherein R^{3a} represents a nitro group and then converting the nitro group into a halogen atom or a group of the abovedefined formula $-NR^4R^5$.

Accordingly, in one particular aspect the present invention provides a process for preparing a compound of formula (IA) which process comprises reacting a compound of the hereinbefore defined formula (II) with a nitrating agent, to provide a compound of formula (IA) wherein R^{3a} represents a nitro group, and thereafter if required carrying out the following optional steps:

- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) converting the nitro group into another group R^{3a} ;
- (iii) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

A compound of formula (II) may be prepared by the dehydrating cyclisation of a compound of formula (III):



(III)


wherein R^{1a} represents R^1 , as defined in relation to formula (IA), or a group convertible to R^1 and R^{2a} represents R^2 , as defined in relation to formula (IA), or a group convertible thereto, A^1 represents $-NO$ or $-NH.CHO$ and A^2 represents $-NH.CH_3$ or $-NH_2$, providing that when A^1 is $-NO$ then A^2 is $-NH.CH_3$ and when

A¹ is -NH.CHO then A² is NH₂; and thereafter, if required, converting any group R^{1a} to R¹ and/or R^{2a} to R². The dehydrating cyclisation of a compound of formula (III) may be carried out under any suitable conditions. Favourably the conditions chosen are these wherein the water formed is removed from the reaction mixture, thus the reaction is generally carried out at an elevated temperature in the range of from 100° C to 200° C, such as in the range of 180° C to 190° C.

In one aspect of the process, especially when A¹ is -NO and A² is -NH.CH₃, the reaction is carried out in a solvent immiscible with water, such as toluene, at the reflux temperature of the solvent, the water being removed using a water-separator.

Suitable values for R^{1a} and R^{2a} include R¹ and R² respectively or nitrogen protecting groups such as benzyl groups.

When R^{1a} or R^{2a} represents other than R¹ or R² respectively, the abovementioned conversions of R^{1a} into R¹ and R^{2a} to R² may be carried out using the appropriate conventional procedure. For example when R^{1a} (or R^{2a}) represents a nitrogen protecting group, such as a benzyl group, the protecting group may be removed using the appropriate conventional procedure, such as catalytic hydrogenation, and the resulting product reacted with a compound of formula (IV):




wherein A and m are as defined in relation to formula (IA) and X represents a leaving group, such as halide, for example bromide or iodide.

The protection of any reactive group or atom, such as the xanthine nitrogen atom may be carried out at any appropriate stage in the aforementioned process. Suitable protecting groups include those used conventionally in the art for the particular group or atom being protected, for example suitable protecting groups for the xanthine nitrogen atoms are benzyl groups.

Protecting groups may be prepared and removed using the appropriate conventional procedure:

For example, N-benzyl protecting groups may be prepared by treating the appropriate compound of formula (II) with benzyl chloride in the presence of a base such as triethylamine. The N-benzyl protecting groups may be removed by catalytic hydrogenation over a suitable catalyst, such as palladium on activated charcoal, in a suitable solvent, such as ethanol conveniently at an elevated temperature, or by treatment with anhydrous aluminium chloride in dry benzene at ambient temperature.

A compound of formula (III) wherein A¹ represents -NH.CHO and R² represents -NH₂ may suitably be prepared from a 6-aminouracil of formula (A) according to the following reaction scheme:




wherein R^{1a} and R^{2a} are as defined in relation to formula (II).

Suitably, the reaction conditions used in the abovementioned reaction scheme are appropriate conven-

tional conditions. In a preferred aspect of the process, the conversion of the 6-aminouracil (A), via (B) and (C), to the corresponding compound of formula (III) and the cyclisation of the compound of formula (III) to the compound of formula (II) are all carried out in-situ, suitably by using an analogous procedure to that of H. Brederick and A. Edenhofer, Chem. Berichte 88, 1306-1312 (1955).

The 6-aminouracils of formula (A) may themselves be prepared by the method of V. Papesch and E. F. Schroder, J. Org. Chem., 16, 1879-90 (1951), or Yozo Ohtsuka, Bull. Chem. Soc. Jap., 1973, 46(2), 506-9.

A compound of formula (III) wherein A¹ represents -NO and A² represents -NH.CH₃ may conveniently be prepared from a 6-chlorouracil of formula (D), according to the following reaction scheme:



wherein R^{1a} and R^{2a} are as defined in relation to formula (II).

Suitably, the reaction conditions used in the last above mentioned scheme are the appropriate conventional conditions, for example those used in the method of H. Goldner, G. Dietz and E. Carstens, Liebigs Annalen der Chemie, 691, 142-158 (1965). The 6-chlorouracil of formula (D) may also be prepared according to the procedure of Dietz et al.

When R^{3a} represent a nitro group, suitable conversions of the nitro group into another group R^{3a} include the following:

- (i) converting the nitro group into a halogen atom;
- (ii) converting the nitro group into an amine group;
- (iii) converting the nitro group into a halogen atom followed by conversion of the halogen atom into a group -NR^{4b}R^{5b} wherein R^{4b} and R^{5b} together with the nitrogen atom to which they attached form an optionally substituted heterocyclic group; and
- (iv) converting the nitro group into an amino group and thereafter alkylating and/or acylating the amino group to provide a group -NR^{4c}R^{5c} wherein R^{4c} represents hydrogen, alkyl or alkylcarbonyl and R^{5c} represents alkyl or alkylcarbonyl.

A nitro group may be converted into a halogen atom by using any convenient halogenating agent.

One suitable halogenating agent is a hydrogen halide, suitably reacted in aqueous conditions for example by using concentration hydrochloric acid or concentrated hydrobromic acid at an elevated temperature, for example in the range of from 50 to 150 °C.

A further suitable halogenating agent is a phosphorous oxyhalide, such as phosphorous oxychloride, which may be reacted in any suitable solvent, such as dimethylformamide, suitably at an elevated temperature for example in the range of from 50 °C to 150 °C.

A nitro group may conveniently be converted into an amino group by conventional reduction methods for example by using tin powder and concentrated hydrochloric acid at ambient temperature or by using sodium dithionite in aqueous methanol at ambient temperature.

When R^{3a} in the compound of formula (IA) represents a halogen atom it may be converted into a group -NR^{4b}R^{5b} by reacting with a reagent of formula (III):



wherein R^{4b} and R^{5b} are as defined above.

The reaction between the compound of formula (IA) and the compound of formula (III) may be carried out in any suitable solvent, such as toluene, at any temperature providing a convenient rate of formation of the product, but suitably at an elevated temperature, such as in the range of from 50° to 180° C, at atmospheric or an elevated pressure.

Suitable alkylation methods for use in the abovementioned conversions include those used conventionally in the art, for example methods using halides, preferably iodides, in the presence of a base such as potassium carbonate in any convenient solvent for example acetonitrile or toluene.

Suitable acylation methods for use in the abovementioned conversions include those used conventionally in the art, thus an amino group may be converted into an alkylcarbonyl amino group by using an appropriate acylating agent, for example an amino group may be converted to an acetylamino group by using acetic anhydride at elevated temperature.

The compounds of formula (I) may be prepared according to the abovementioned methods or, as appropriate, by the methods of the abovementioned publications.

The active compound may be formulated for administration by any suitable route, the preferred route depending upon the disorder for which treatment is required, and is preferably in unit dosage form or in a form that a human patient may administer to himself in a single dosage. Advantageously, the composition is suitable for oral, rectal, topical, parenteral, intravenous or intramuscular administration or through the respiratory tract. Preparations may be designed to give slow release of the active ingredient.

The compositions of the invention may be in the form of tablets, capsules, sachets, vials, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations such as oral or sterile parenteral solutions or suspensions. Topical formulations are also envisaged where appropriate.

In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit dose.

Unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate; disintegrants, for example starch, polyvinylpyrrolidone, sodium starch glycollate or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulphate.

The solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers.

Such operations are of course conventional in the art. The tablets may be coated according to methods well known in normal pharmaceutical practice, in particular with an enteric coating.

Oral liquid preparations may be in the form of, for example, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel, hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid; and if desired conventional flavouring or colouring agents.

Compositions may also suitably be presented for administration to the respiratory tract as a snuff or an aerosol or solution for a nebulizer, or as a microfine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case the particles of active compound suitably have diameters of less than 50 microns, such as from 0.1 to 50 microns, preferably less than 10 microns, for example from 1 to 10 microns, 1 to 5 microns or from 2 to 5 microns. Where appropriate, small amounts of other anti-asthmatics and bronchodilators, for example sympathomimetic amines such as isoprenaline, isoetharine, salbutamol, phenylephrine and ephedrine; xanthine derivatives such as theophylline and aminophylline and corticosteroids such as prednisolone and adrenal stimulants such as ACTH may be included.

For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, and, depending on the concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, a preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead

of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The compositions may contain from 0.1% to 99% by weight, preferably from 10-60% by weight, of the active material, depending on the method of administration.

Compounds of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, may also be administered as a topical formulation in combination with conventional topical excipients.

Topical formulations may be presented as, for instance, ointments, creams or lotions, impregnated dressings, gels, gel sticks, spray and aerosols, and may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams. The formulations may contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions.

Suitable cream, lotion, gel, stick, ointment, spray or aerosol formulations that may be used for compounds of formula (I) or if appropriate a pharmaceutically acceptable salt thereof, are conventional formulations well known in the art, for example, as described in standard text books of pharmaceuticals and cosmetics, such as *Harry's Cosmeticology* published by Leonard Hill Books, *Remington's Pharmaceutical Sciences*, and the *British and US Pharmacopoeias*.

Suitably, the compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, will comprise from about 0.5 to 20% by weight of the formulation, favourably from about 1 to 10%, for example 2 to 5%.

The dose of the compound used in the treatment of the invention will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and the relative efficacy of the compound. However, as a general guide suitable unit doses may be 0.1 to 1000mg, such as 0.5 to 200, 0.5 to 100 or 0.5 to 10 mg, for example 0.5, 1, 2, 3, 4 or 5 mg; and such unit doses may be administered more than once a day, for example 2, 3, 4, 5 or 6 times a day, but preferably 1 or 2 times per day, so that the total daily dosage for a 70kg adult is in the range of about 0.1 to 1000 mg, that is in the range of about 0.001 to 20 mg/kg/day, such as 0.007 to 3, 0.007 to 1.4, 0.007 to 0.14 or 0.01 to 0.5 mg/kg/day, for example 0.01, 0.02, 0.04, 0.05, 0.06, 0.08, 0.1 or 0.2 mg/kg/day; and such therapy may extend for a number of weeks or months.

When used herein the term 'pharmaceutically acceptable' encompasses materials suitable for both human and veterinary use.

No toxicological effects have been established for the compounds of formula (I) in the abovementioned dosage ranges.

The following pharmacological data and examples illustrate the invention. The following preparations illustrate the preparation of intermediates to the novel compounds of formula (IA).

Example 1

1,3-Di-n-butyl-8-nitro xanthine

1,3-Di-n-butylxanthine (73g, 0.28mol) was dissolved in acetic acid (120ml) and then treated with concentrated nitric acid (49g) at 87°C. After 1 hour, the mixture was cooled to 5°C, the resulting yellow precipitate filtered off and washed with water (50ml). The yellow crystals were dissolved in dichloromethane and washed twice with water. The separated organic layer was then dried (anhydrous sodium sulphate) and concentrated to give a crystalline product, yield 73g (86%), m.pt 168°C

¹H NMR (CDCl₃/DMSO):

ppm: 0.93 (t, J=6.3Hz, 6H), 1.1-2.0 (m, 8H), 3.8-4.25 (m, 4H).

Example 2

1,3-Di-cyclopropylmethyl-8-nitro xanthine

1,3-Di-cyclopropylmethyl xanthine (20g, 0.076mol) was dissolved in acetic acid (33ml) and then treated with concentrated nitric acid (13.2g) at 87°C. After 1 hour, the mixture was cooled to 5°C and the resulting yellow precipitate filtered off. The yellow crystals were dissolved in dichloromethane and washed with water.









The separated organic layer was then dried over anhydrous sodium sulphate and concentrated in vacuo. The product crystallized from the concentrate to yield a yellow crystalline product yield 12.2g, (56.5%), m.pt. 207 °C (with decomposition).

¹H NMR (CDCl₃):

ppm: 0.35-0.7 (m, 8H), 1.1 -1.7 (m, 2H), 3.95-4.2 (m, 4H), 9.0-11.0 (br exchanges with D₂O, 1H).

The following compounds were prepared using an analogous procedure to that described in Example 1. The appropriate 1,3-di-cycloalkylmethyl xanthine substrates were prepared according to the procedures described herein in and in United Kingdom Patent Application No. 8826595.4.

Table 1

Ex. No	R ¹	R ²	R ³	M.pt (°C)	¹ H NMR Spectrum: (CDCl ₃ or CDCl ₃ /DMSO, ppm)
3			NO ₂	220	1.7-2.2 (m, 12H) 2.5-3.1 (m, 2H) 4.1-4.3 (m, 4H)
4			NO ₂	148-150	1.1 -2.0 (m, 16H) 2.15-2.7 (m, 2H) 4.15 (d, J=7.7Hz, 4H)
5			NO ₂	140	0.75-2.2 (m, 22H) 3.7 -4.1 (m, 4H)
6			NO ₂	>250	0.7 -1.4 (m, 8H) 2.5 -3.4 (m, 2H)

Example 7

1,3-Di-n-butyl-8-amino xanthine hydrochloride

1,3-Di-n-butyl-8-nitro xanthine from Example 1 (8.5g) was suspended in concentrated hydrochloric acid (85ml) and then treated at room temperature with tin powder (14.5g) in small portions. After stirring for 10 minutes the yellow colour of the suspension disappeared. Thereafter the precipitate was filtered off and recrystallised twice from ethanol. The product formed colourless crystals, yield 5.5g (63%) m.pt>250 °C

¹H NMR (DMSO):

ppm: 0.90 (t, J=6.1Hz, 6H), 1.05-1.9 (m, 8H), 3.65-4.15 (m, 4H), 6.9 (br, exchanges with D₂O, 4H).

Example 8

1,3-Di-n-butyl-8-amino xanthine

Neutralisation of the hydrochloride from Example 6 with 1N sodium hydroxide solution gave the 1,3-di-

n-butyl-8-amino xanthine as a crystalline compound, yield 92%, m.pt. 89° C.

Example 9

1,3-Di-cyclopropylmethyl-8-amino xanthine

1,3-Di-cyclopropylmethyl-8-nitro xanthine (4g, 0.014mol), suspended in 50ml of concentrated hydrochloric acid, was treated with small portions of tin (8g) at room temperature. The mixture was then stirred at room temperature for two hours.

The resulting precipitate was filtered off and crystallised from ethanol to give white crystals of the title product, yield 0.9g (23%) m.pt. 281° C.

In an alternative procedure, using sodium dithionite as reducing agent (in methanol-water mixture). The yield was 36% (compare Example 13).







¹H NMR (CDCl₃):

ppm: 0.3-0.6 (m, 8H), 1.0-1.6 (m, 2H), 3.7-4.0 (m, 4H), 5.75 (br, 2H), 10.84 (br. exchanges with D₂O, 1H).

Examples 10 to 12

The following compounds were prepared using an analogous procedure to that described above for the preparation of a compound of Example 7.

Table 2

Ex. No	R ¹	R ²	R ³	M.pt (°C)	¹ H NMR Spectrum: (CDCl ₃ or CDCl ₃ /DMSO, ppm)
10			NH ₂	124 (decomp)	1.65-2.2 (m, 12H) 2.5-3.1 (m, 2H) 3.85-4.2 (m, 4H) 6.5-8.5 (br, 4H)
11			NH ₂	300	1-1.9 (m, 16H) 2.15-2.7 (m, 2H) 3.65-4.0 (m, 4H) 6.45 (br, 2H) 11.20 (br, 1H)
12			NH ₂	300	0.7-2.2 (m, 22H) 3.65-3.95 (m, 4H) 6.51 (s, 2H) 10-13 (br, 1H)

Example 13

1,3-Di-cyclopropyl-8-amino xanthine

1,3-Di-cyclopropyl-8-nitro xanthine (0.4g, 0.0014mol) was dissolved in methanol (20ml) and treated with a sodium dithionite solution in water (0.5g in 5 ml) at room temperature with stirring. After stirring for three

hours the solvent was removed in vacuo, the residue taken up with dichloromethane and extracted with water (40ml).

After drying the organic layer over anhydrous sodium sulphate, the solvent was removed and the residue crystallised from ethanol, yield 0.06g (17%), m.pt. >250 °C.

5 ¹H NMR (CDCl₃):

ppm: 0.6-1.4 (m, 8H), 2.6-3.25 (m, 2H), 7.8 (s, 1H), 12 (br. exchanges with D₂O, 1H).

Example 14

10

1,3-Di-n-butyl-8-acetamido xanthine

1,3-Di-n-butyl-8-amino xanthine (0.5g), hydrochloride in toluene (30ml) was stirred for 30 minutes with triethylamine (0.16g). After addition of acetic anhydride (0.32g), the mixture was refluxed for 6 hours. The reaction mixture was extracted with water (4 x 30ml), the organic layer separated and dried over anhydrous sodium sulphate. The solvent was then evaporated to yield the product, yield 0.1g (20%), mpt. 180 °C.

¹H NMR (CDCl₃):

ppm: 0.93 (t, J=6.4Hz, 6H), 1.1-1.9 (m, 8H), 2.27 (s, 3H), 4.01 (t, J=6.7Hz), 8.9 (br, exchanges with D₂O, 1H).

20

Example 15

25

1,3-Di-n-butyl-8-chloro xanthine

1,3-Di-n-butyl-8-nitro xanthine (0.5g, 0.0016mol) was refluxed for 18 hours with concentrated hydrochloric acid (8ml). The reaction mixture was extracted with dichloromethane (20ml), the organic layer washed with water to neutrality and then dried over anhydrous sodium sulphate. The solvent was then removed by evaporation in vacuo and the residue was recrystallised from ethanol, to give the title compound, yield 0.38g (73%), m.pt. 152 °C.

¹H NMR (CDCl₃):

ppm: 0.97 (t, J=6.1 Hz, 6H), 1.1-2.0 (m, 8H), 4.11 (t, J=7Hz, 4H), 13.1 (br., exchanges with D₂O, 1H).

35

Example 16

1,3-Di-n-butyl-8-bromo xanthine

1,3-Di-n-butyl-8-bromo xanthine was prepared from 1,3-di-n-butyl-8-nitro xanthine (0.5g, 0.0016mol) and concentrated hydrobromic acid (8ml) using the procedure as described in Example 15. The title product was obtained after recrystallisation from ethanol, yield 0.4g (91%), m.pt. 178 °C.

45 ¹H NMR (CDCl₃):

ppm: 0.97 (t, J=6.1Hz, 6H), 1.1-2.0 (m, 8H), 4.11 (t, J=6.9Hz, 4H), 13.3 (br. exchanges with D₂O, 1H).

Example 17

50

1,3-Di-cyclopropylmethyl-8-chloro xanthine

1,3-Di-cyclopropylmethyl-8-nitro xanthine (6g, 0.023mol) was dissolved in dimethylformamide (20ml) and reacted with phosphorous oxychloride (14g) for 1 hour at 120 °C. The mixture was then treated with water and stirred for 1 hour at room temperature. The precipitate was filtered off, dissolved in ethyl acetate, dried over anhydrous sodium sulphate and the solvent was removed in vacuo, yield 2.5g (40%), m.pt. 220 °C.

¹H NMR (CDCl₃):

ppm: 0.35-0.65 (m, 8H), 1.05-1.65 (m, 2H), 4.1 (d, J=7.1 Hz, 4H), 13.4 (br. exchanges with D₂O, 1H).

5 Example 18

1,3-Di-cyclohexyl-8-chloro xanthine

10 1,3-Di-cyclohexyl-8-chloro xanthine was prepared from 1,3-di-cyclohexyl-8-nitro xanthine (2g, 0.006mol) and phosphorous oxychloride (3.9g) in dimethylformamide (6ml), using an analogous procedure to that described in Example 17. The product was obtained as a crystalline product after recrystallisation from ethyl acetate, m.pt., 135 °C.

¹H NMR (CDCl₃):

15 ppm: 1.0-2.7 (m, 20H), 4.3-5.0 (m, 2H).

Example 19

20

1,3-Di-n-butyl-8-piperidino xanthine

1,3-Di-n-butyl-8-bromo xanthine (2g, 0.0029mol) was dissolved in toluene (50ml). After addition of piperidine (5g, 0.0058mol) the mixture was refluxed for 9 hours. The reaction mixture was then extracted with water (4 x 30ml), the organic layer dried over anhydrous sodium sulphate and the solvent removed in vacuo. The residue was recrystallised from ethanol to give the title product, yield 0.4g (20%), m.pt. 221 °C.

¹H NMR (CDCl₃):

ppm: 0.85-1.15 (m, 6H), 1.15-2.1 (m, 14H), 3.5-3.8 (m, 4H), 11 (br. exchanges with D₂O, 1H).

30

Example 20

1,3-Di-cyclopropylmethyl-8-morpholino xanthine

35

1,3-Di-cyclopropylmethyl-8-morpholino xanthine was prepared from 1,3-di-cyclopropylmethyl-8-chloroxanthine (0.3g, 0.001mol) and morpholine (0.2g, 0.0022mol) using an analogous procedure to that described in Example 19. The title product was obtained as a crystalline solid, m.pt. >250 °C, yield 0.09g (26%).

¹H NMR (CDCl₃):

40 ppm: 0.3-0.65 (m, 8H), 1.0-1.7 (m, 2H), 3.5-4.2 (m, 12H), 11.4 (br. exchanges with D₂O, 1H).

Example 21

45

1,3-Di-n-butyl-8-pyrrolidinyl xanthine

The title compound was prepared from 1,3-di-n-butyl-8-bromo xanthine (1g, 0.0029mol) and pyrrolidine (0.041g, 0.0057mol) using an analogous procedure to that described in Example 19. The title product was obtained as a crystalline solid, m.pt. >250 °C.

¹H NMR (CDCl₃):

ppm: 0.097 (t, J=6.3Hz, 6H), 1.1-2.25 (m, 12H), 3.5-3.8 (m, 4H), 3.8-4.2 (m, 4H), 10.9 (br. exchanges with D₂O, 1H).

55

Example 22

1,3-Di-cyclopropylmethyl-8-pyrrolidinyl xanthine

The title compound was prepared from 1,3-di-cyclopropylmethyl-8-chloro xanthine (0.3g, 0.0011mol) and pyrrolidine (0.2g, 0.0028mol) using an analogous procedure to that described in Example 19. The title product was obtained as a crystalline solid, m.pt. >250 °C.

¹H NMR (CDCl₃):

ppm: 0.3-0.65 (m, 8H), 1.1-1.8 (m, 2H), 1.9-2.2 (m, 4H), 3.5-3.8 (m, 4H), 3.8-4.1 (m, 4H) 10.6 (br. exchanges with D₂O, 1H).

Example 231,3-Di-cyclopropylmethyl-8-piperidinyl xanthine

The title compound was prepared from 1,3-di-cyclopropylmethyl-8-bromo xanthine (1.2g, 0.0037mol) and piperidine (0.79g, 0.009mol) using an analogous procedure to that described in Example 19. The title product was obtained as a crystalline solid, m.pt. >250 °C.

¹H NMR (CDCl₃):

ppm: 0.3-0.6 (m, 8H), 1.05-1.55 (m, 2H), 1.55-1.9, (m, 6H), 3.45-3.8 (m, 4H), 3.8-4.05 (m, 4H), 13.3 (br., exchanges with D₂O, 1H).

Example 241,3-Di-cyclohexylmethyl-8-piperidinyl xanthine

The title compound was prepared from 1,3-di-cyclohexylmethyl-8-bromo xanthine (0.7g, 0.0017mol) and piperidine (0.28g, 0.003mol) using an analogous procedure to that described in Example 19. The title product was obtained as a crystalline solid, m.pt. 266 °C.

¹H NMR (CDCl₃):

ppm: 0.75-2.2 (m, 28H), 3.5-3.75 (m, 4H), 3.75-4.05 (m, 4H), 10.72 (br. exchanges with D₂O, 1H).

Example 251,3-Di-cyclohexylmethyl-8-bromo xanthine

The title compound was prepared from 1,3-di-cyclohexylmethyl-8-nitro xanthine (1g, 0.0026mol) and concentrated hydrobromic acid (40ml, 48%) over 32 hours using an analogous procedure to that described in Example 15. The title product was obtained as a crystalline solid, m.pt. 247 °C.

¹H NMR (CDCl₃):

ppm: 0.75-2.2 (m, 22H), 3.89 (d, J = 7.2Hz, 4H), 13.45 (br. exchanges with D₂O, 1H).

Example 261,3-Di-cyclohexyl-8-nitro xanthine

The title compound was prepared from 1,3-di-cyclohexyl xanthine (1.5g, 0.0044mol) concentrated nitric acid (0.56g) and acetic acid (1.9ml) using an analogous procedure to that described in Example 1. The title product was obtained as a crystalline solid, m.pt. >250 °C.

¹H NMR (CDCl₃):

ppm: 0.8-2.7 (m, 20H), 4-5 (m, 2H).

Preparation 11,3-Di-cyclopropylmethyl xanthine

1,3-Di-cyclopropylmethyl xanthine was prepared using an analogous procedure to that described in Chem. Berichte 88, 1306-1312, 1955: 20.2g (0.0855 mol) of 1,3-di-cyclopropylmethyl-6-amino-uracil was dissolved in 100ml of formamide, then 5.9g sodium nitrite was added and at 60 °C 13.4ml formic acid was added slowly with stirring. After the colour had changed from yellow to violet, the mixture was heated up to 100 °C and 3.1g of sodium dithionite (Na₂S₂O₄) was added in small portions.

Then the mixture was heated to 180-190 °C and held at this temperature for 30 minutes.

After cooling, the precipitate was sucked off, washed with 50ml of water and recrystallised from toluene. Yield: 22.5g, m.p. 203 °C.

¹H NMR (CDCl₃):

ppm: 0.44-0.54 (8H, q); 1.18-1.57 (2H, m); 3.98-4.12 (4H, 2d); 7.81 (1H, s); 12.8-13.2 (1H, s, exch. with D₂O).

Preparation 21,3-Di-cyclobutylmethyl xanthine

1,3-Di-cyclobutylmethyl xanthine was prepared from 1,3-dicyclobutylmethyl-6-aminouracil using an analogous procedure to that described in Preparation 1. The title compound was isolated as a crystalline solid, m.p. 191 °C.

¹H NMR (CDCl₃):

ppm: 1.6-2.3 (12H, m); 2.4-3.2 (2H, m); 4.16 (2H, d, J=7.0Hz); 4.21 (2H, d, J=7.3Hz); 7.76 (1H, d, J=1.3Hz, exch. with D₂O to give s) 12.7 (1H, br.s, exch. with D₂O).

Preparation 31,3-Di-cyclopentylmethyl xanthine

1,3-Di-cyclopentylmethyl xanthine was prepared from 1,3-di-cyclopentylmethyl-6-aminouracil using an analogous procedure to that described in Preparation 1. The title compound was isolated as a crystalline solid, m.p. 208 °C.

¹H NMR (CDCl₃):

ppm: 1.0-2.0 (16H, m); 2.2-2.9 (2H, m); 4.0-4.3 (4H, m); 7.78 (1H, d, J=1.2Hz, exch. with D₂O to give s); 12.9 (1H, br.s, exch. with D₂O).

Preparation 41,3-Di-cyclohexylmethyl xanthine

1,3-Di-cyclohexylmethyl xanthine was prepared from 1,3-di-cyclohexylmethyl-6-aminouracil using an analogous procedure to that described in Preparation 1. The title compound was isolated as a crystalline solid, m.p. 237 °C.

¹H NMR (CDCl₃):

ppm: 0.8-2.2 (22H, m); 3.85-4.15 (4H, m (dd)); 7.73 (1H, s); 13.1 (1H, br.s, exch. with D₂O).

Preparation 5

1,3-Di-cyclopropylmethyl-6-aminouracil

1,3-Di-cyclopropylmethyl-6-aminouracil was prepared using an analogous procedure to that described in J. Org. Chem. 16, 1879-1890, (1951):

22.6g (0.138mol) of the N,N'-dicyclopropylmethyl-urea (from Preparation 1) was treated with 44ml (0.43mol) of acetic anhydride and 14g (0.165mol) of cyanocetic-acid at 70 °C for 2 hours.

After cooling and the addition of 15ml of water, 40ml of 50% NaOH/water-solution was dropped slowly onto the mixture at 45 °C with stirring.

After stirring for 1 hour at room temperature, the strongly alkaline solution was separated and the oily residue washed carefully with 60ml water.

The semi-solid residue was dissolved in 220ml methanol and dropped into 1 litre of water with stirring. Thereby the product crystallised. Yield: 25.5g, 78.5% approx., m.p. 85-95 °C (wax-like).

Preparation 61,3-Di-cyclopentylmethyl-6-aminouracil

1,3-Di-cyclopentylmethyl-6-aminouracil was prepared from N,N'-di-cyclopentylmethyl urea using a procedure analogous to that described in Preparation 5. The title compound was isolated as a crystalline solid, m.p. 108 °C.

¹H NMR (CDCl₃):

ppm: 1.0-2.6 (18H, m); 3.86 (4H, d, J = 7.4Hz); 4.98 (3H, m, 2H exch. with D₂O).

Preparation 71,3-Di-cyclohexylmethyl-6-aminouracil

1,3-Di-cyclohexylmethyl-6-aminouracil was prepared from N,N'-di-cyclohexylmethyl urea using a procedure analogous to that described in Preparation 5. The title compound was isolated as a crystalline solid, m.p. 185 °C.

Preparation 8N,N'-Di-cyclopropylmethyl urea

N,N'-Di-cyclopropylmethyl urea, m.p. 124 °C, was prepared using a procedure analogous to that described in J. Org. Chem. 16, 1879-1890, (1951):

68.2g (0.634mol) cyclopropylmethylamine-hydrochloride in 800ml of water was treated with 25g sodium hydroxide dissolved in 100ml of water and the mixture cooled to -15 °C.

Phosgene, 33g was then slowly introduced through a capillary tube with stirring. Thereafter the mixture was stirred for 1 hour and, as necessary, after acidification with 0.1 N HCl, the product was extracted with dichloromethane.

After washing with water and drying over anhydrous sodium sulphate the product was obtained after evaporation of the solvent. Yield: 21g, 40% approx.

From the aqueous phase, 20g of the unreacted adduct (cyclopropylmethylamine-hydrochloride) can be obtained.

¹H NMR (CDCl₃):

ppm: 0.06-0.59 (8H, m); 0.72-1.06 (2H, m); 3.01-3.09 (4H, d); 4.66 (1H, br.s, exch. with D₂O).

Preparation 9

N,N'-Dicyclobutylmethyl urea

N,N'-Di-cyclobutylmethyl urea was prepared from cyclobutylmethylamine using a procedure analogous to that described in Preparation 8. The title compound was isolated as a crystalline solid, m.p. 155° C.

¹H NMR (CDCl₃):

ppm: 1.4-2.8 (14H, m); 3.0-3.3 (4H, m); 4.59 (2H, br.s, exch. with D₂O).

Preparation 10N,N'-Di-cyclopentylmethyl urea

N,N'-Di-cyclopentylmethyl urea was prepared from cyclopentylmethylamine using a procedure analogous to that described in Preparation 8. The title compound was isolated as a crystalline solid, m.p. 150° C.

¹H NMR (CDCl₃):

ppm: 1.0-2.2 (18H, m); 2.9-3.2 (4H, m); 4.59 (2H, br.s, exch. with D₂O).

Preparation 11N,N'-Di-cyclohexylmethyl urea

N,N'-Di-cyclohexylmethyl urea was prepared from cyclohexylmethylamine using a procedure analogous to that described in Preparation 8. The title compound was isolated as a crystalline solid, m.p. 159° C.

PHARMACOLOGICAL DATAa) Inhibition of Cyclic AMP PhosphodiesteraseProcedure

The procedure used was that described by Arch, J.R.S. and Newsholme, E.A. in Biochem. J. 158, 603, (1976):

Erythrocytes were obtained from Na-citrate (16 mM; 0.1 ml/ml blood) anticoagulated blood by repeated centrifugation with removal of the buff coat and washing with an isotonic buffer (composition in mM: NaCl 13.7, KCl 4, CaCl₂.2 H₂O 1.8, Na₂HPO₄.12 H₂O 0.8, NaH₂PO₄ 0.2, MgSO₄.7 H₂O 0.7, Hepes 3.4; pH 7.4).

The phosphodiesterase was extracted by mixing the erythrocytes with 4 volumes of 7 mM phosphate buffer, pH 7.4, followed by sonification (3 x 10 sec; 100 W) and then centrifuging for 30 min at 4200 x g.

All supernatants were diluted in the extraction medium and assayed for phosphodiesterase activity within 6 hours of preparation, using the radiochemical procedure described in the above mentioned reference.

Results		
	Example 1	Ki [μ M] c-AMP phosphodiesterase (erythrocytes)
	No.	
5	1	17
	2	15.9
10	3	6.1
	4	4.8
	5	5.4
	7	1.3
	9	1.6
15	10	0.53
	11	0.57
	13	14
	17	15.1
	18	23
20	19	<100
	20	11.5
	22	29.7
	23	55.9
25	25	7.9

b) Induction of blood eosinophilia and the effects of drugs.

Animals

Male Charles River Sprague Dawley rats weighing between 250 to 300g were used.

The method used was a modification of that described by Laycock et al (Int. Arch. Appl. Immunol, (1986), 81, 363).

Sephadex G200, particle size 40 to 120 micron, was suspended in isotonic saline at 0.5mg/ml, and stored for 48h at 4 °C. 1ml of the suspension was given intravenously to rats on days 0,2 and 5. A control group received saline. The test compound was given before the Sephadex on each occasion, with a contact time expected to give maximum activity at the time of the Sephadex administration. Blood was taken from the tail vein of the rats on day 7 for the determination of total and differential leucocyte counts.

A control group of at least 6 animals was included each time a compound was evaluated. The control group received Sephadex and the vehicle without test compound. The results in the drug treated animals were compared with the control group. Alternatively, if the mean for the control group for any experiment was not statistically different from the mean of the sum of all of the control groups, then the treated animal results for that experiment were compared with the mean of the sum of all the control groups.

Total and differential leucocyte counts.

20 μ l samples of blood, taken from the tail vein of the rats, were added to 10ml of Isoton II and, within 30min, Zaponin (3 drops) was added, to lyse the erythrocytes. Five minutes later the total cell count was determined using a Coulter Counter Model DN. Differential leucocyte counts were carried out by fixing and staining a blood smear on a microscopic slide with May-Grunwald and Giemsa stains. A minimum of 400 cells were counted on each slide.

Statistics

Probability values were calculated using the Student's t test.

Results

The effect of the test compound upon Sephadex induced eosinophilia in the rat is set out in Table 3. The test compound was given orally 30 minutes before each injection of Sephadex. The results indicate that the test compound inhibits the induced eosinophilia in a dose dependent manner.

Table 3


Test Compound	Dose mg/kg (orally - 30 mins)	% of Control Mean \pm SEM (n = 12 or more)
Vehicle dosed control + sephadex i.v.	-	100 \pm 7
Negative control saline i.v.	-	10 \pm 0.9 ^{***}
Example 1	25.0	109 \pm 6
Example 7	25.0	91 \pm 6
Example 9	2.0	48 \pm 7 ^{***}
	0.2	61 \pm 8 ^{**}
Example 10	5.0	47 \pm 4 ^{***}
Notes		

^{**} p < 0.01

^{***} p < 0.001

Claims

1. The use of a compound of formula (I):



(I)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R¹ and R² each independently represent alkyl or a moiety of formula (a):

$-(CH_2)_m-A$ (a)


wherein m represents zero or an integer 1, 2 or 3;

A represents a substituted or unsubstituted cyclic hydrocarbon radical; and

R³ represents a halogen atom, a nitro group, or a group $-NR^4R^5$ wherein R⁴ and R⁵ each independently represents hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are attached forming an optionally substituted, heterocyclic group; for the manufacture of a medicament for the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischaemic events and/or peripheral vascular disease and/or proliferative skin diseases and/or disorders of the respiratory tract and/or the treatment or prophylaxis of disorders associated with increased numbers of eosinophils and allergic disorders associated with atopy.

2. A pharmaceutical composition comprising a compound of formula (I), or if appropriate a pharmaceutically acceptable salt thereof, providing that in the compound of formula (I) when R¹ and R² both represent methyl then R³ is not a nitro group and a pharmaceutically acceptable carrier therefor.

3. A compound of formula (IA):



(IA)

or if appropriate a pharmaceutically acceptable salt thereof, characterised in that:

R¹ and R² each independently represent alkyl or a moiety of formula (a):

-(CH₂)_m-A (a)

wherein m represents zero or an integer 1, 2 or 3; A represents a substituted or unsubstituted cyclic hydrocarbon radical, providing that when R¹ represents methyl then R² is not methyl; and R^{3a} represents a halogen atom, a nitro group, or a group -NR⁴R⁵ wherein R⁴ and R⁵ each independently represent hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are attached forming an optionally substituted heterocyclic group.

4. A compound according to claim 3, wherein R¹ represents a moiety of formula (a).

5. A compound according to claim 3 or claim 4, wherein R² represents a moiety of formula (a).

6. A compound according to any one of claims 3 to 5, wherein R¹ and R² each independently represent a moiety of formula (a).

7. A compound according to any one of claims 3 to 6, wherein A represents a substituted or unsubstituted C₃₋₈ cycloalkyl group.

8. A compound according to any one of claims 3 to 7, wherein A represents a substituted or unsubstituted cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl group.

9. A compound according to any one of claims 3 to 8, wherein A represents a cyclopropyl group or a cyclobutyl group.

10. A compound according to any one of claims 3 to 9, wherein A represents a cyclopropyl group.

11. A compound according to any one of claims 3 to 10, wherein R^{3a} is a nitro group or a group -NHR⁴ wherein R⁴ represents hydrogen or alkylcarbonyl.

12. A compound according to any one of claims 3 to 11, wherein R^{3a} represents an amino group.

13. A compound according to any one of claims 3 to 10, wherein R^{3a} represents a halogen atom.

14. A compound according to any one of claims 3 to 10, wherein R^{3a} represents a pyrrolidinyl, piperidinyl or morpholinyl group.


15. A compound according to claim 3 selected from the group consisting of:

- 1,3-di-n-butyl-8-nitro xanthine;
- 1,3-di-cyclopropylmethyl-8-nitro xanthine;
- 1,3-di-cyclobutylmethyl-8-nitro xanthine;
- 1,3-di-cyclopentylmethyl-8-nitro xanthine;
- 1,3-di-cyclohexylmethyl-8-nitro xanthine;
- 1,3-di-n-butyl-8-amino xanthine;
- 1,3-di-cyclopropylmethyl-8-amino xanthine;
- 1,3-dicyclobutylmethyl-8-amino xanthine;
- 1,3-di-cyclopentylmethyl-8-amino xanthine;
- 1,3-di-cyclohexylmethyl-8-amino xanthine;
- 1,3-di-cyclopropyl-8-amino xanthine;
- 1,3-di-n-butyl-8-acetamido xanthine;
- 1,3-di-n-butyl-8-chloro xanthine;
- 1,3-di-n-butyl-8-bromo xanthine;
- 1,3-di-cyclopropylmethyl-8-chloro xanthine;
- 1,3-di-cyclohexyl-8-chloro xanthine;

1,3-di-n-butyl-8-piperidino xanthine;
 1,3-di-cyclopropylmethyl-8-morpholino xanthine;
 1,3-di-n-butyl-8-pyrrolidinyl xanthine;
 1,3-di-cyclopropylmethyl-8-pyrrolidinyl xanthine;
 1,3-di-cyclopropylmethyl-8-piperidinyl xanthine;
 1,3-di-cyclohexylmethyl-8-piperidinyl xanthine;
 1,3-di-cyclohexylmethyl-8-bromo xanthine; and
 1,3-di-cyclohexyl-8-nitro xanthine; or if appropriate a pharmaceutically acceptable salt thereof.

16. A compound according to claim 3, being 1,3-di-cyclopropylmethyl-8-amino xanthine or a pharmaceutically acceptable salt thereof.

17. A process for the preparation of a compound of formula (IA), characterised in that the process comprises reacting a compound of formula (II):



(II)

wherein R^{1a} represents R^1 , as defined in relation to formula (IA), or a group convertible to R^1 and R^{2a} represents R^2 , as defined in relation to formula (IA), or a group convertible thereto, with a reagent capable of substituting the C-8 hydrogen of the compound of formula (II) with a group R^{3b} wherein R^{3b} represents R^{3a} , as defined above in relation to formula (IA), or a group convertible thereto; and thereafter, if required carrying out one or more of the following optional steps:

- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) when R^{3b} is not R^{3a} , converting R^{3b} to R^{3a} ;
- (iii) converting a compound of formula (IA) into a further compound of formula (IA);
- (iv) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

18. A process according to claim 17, which process comprises reacting a compound of formula (II) with a nitrating agent, to provide a compound of formula (IA) wherein R^{3a} represents a nitro group, and thereafter if required carrying out the following optional steps:


- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) converting the nitro group into another group R^{3a} ;
- (iii) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

19. A process according to claim 18 wherein the conversion of the nitro group into another group R^{3a} is selected from the list consisting of:

- (i) converting the nitro group into a halogen atom;
- (ii) converting the nitro groups into an amine group;
- (iii) converting the nitro group into a halogen atom followed by conversion of the halogen atom into a group $-NR^{4b}R^{5b}$ wherein R^{4b} and R^{5b} together with the nitrogen atom to which they attached form an optionally substituted, heterocyclic group; and
- (iv) converting the nitro group into an amino group and thereafter alkylating and/or acylating the amino group to provide a group $-NR^{4c}R^{5c}$ wherein R^{4c} represents hydrogen, alkyl or alkylcarbonyl and R^{5c} represents alkyl or alkylcarbonyl.


Claims for the following Contracting State: ES

1. A process for preparing a compound of formula (IA):




(IA)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R^1 and R^2 each independently represent alkyl or a moiety of formula (a):



wherein m represents zero or an integer 1, 2 or 3, A represents a substituted or unsubstituted cyclic hydrocarbon radical, providing that when R^1 represents methyl then R^2 is not methyl; and R^{3a} represents a halogen atom, a nitro group, or a group $-NR^4R^5$ wherein R^4 and R^5 each independently represent hydrogen, alkyl or alkylcarbonyl or R^4 and R^5 together with the nitrogen to which they are attached forming an optionally substituted heterocyclic group; which process comprises reacting a compound of formula (II):



(II)

wherein R^{1a} represents R^1 , as defined in relation to formula (IA), or a group convertible to R^1 and R^{2a} represents R^2 , as defined in relation to formula (IA), or a group convertible thereto, with a reagent capable of substituting the C-8 hydrogen of the compound of formula (II) with a group R^{3b} wherein R^{3b} represents R^{3a} , as defined above in relation to formula (IA), or a group convertible thereto; and thereafter, if required carrying out one or more of the following optional steps:

- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) when R^{3b} is not R^{3a} , converting R^{3b} to R^{3a} ;
- (iii) converting a compound of formula (IA) into a further compound of formula (IA);
- (iv) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

2. A process according to claim 1, for preparing a compound wherein R^1 represents a moiety of formula (a).

3. A process according to claim 1 or claim 2, for preparing a compound wherein R^2 represents a moiety of formula (a).

4. A process according to any one of claims 1 to 3, for preparing a compound wherein R^1 and R^2 each independently represent a moiety of formula (a).

5. A process according to any one of claims 1 to 4, for preparing a compound wherein A represents a substituted or unsubstituted C_3-8 cycloalkyl group.

6. A process according to any one of claims 1 to 5, for preparing a compound wherein A represents a substituted or unsubstituted cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl group.

7. A process according to any one of claims 1 to 6, for preparing a compound wherein A represents a cyclopropyl group or a cyclobutyl group.

8. A process according to any one of claims 1 to 7, for preparing a compound wherein A represents a cyclopropyl group.

9. A process according to any one of claims 1 to 8, for preparing a compound wherein R^{3a} is a nitro group or a group $-NHR^4$ wherein R^4 represents hydrogen or alkylcarbonyl.

10. A process according to any one of claims 1 to 9, for preparing a compound wherein R^{3a} represents

an amino group.

11. A process according to any one of claims 1 to 8, for preparing a compound wherein R^{3a} represents a halogen atom.

12. A process according to any one of claims 1 to 8, for preparing a compound wherein R^{3a} represents a saturated heterocyclic group having single or fused rings, each ring having 5 to 7 ring atoms which ring atoms optionally comprise up to two additional heteroatoms selected from O, N or S.

13. A process according to claim 12, for preparing a compound wherein R^{3a} represents a pyrrolidinyl, piperidinyl or morpholinyl group.

14. A process according to any one of claims 1 to 13, for preparing a compound wherein m represents

1. 15. A process according to claim 1 for preparing a compound selected from the group consisting of:

- 1,3-di-n-butyl-8-nitro xanthine;
- 1,3-di-cyclopropylmethyl-8-nitro xanthine;
- 1,3-di-cyclobutylmethyl-8-nitro xanthine;
- 1,3-di-cyclopentylmethyl-8-nitro xanthine;
- 1,3-di-cyclohexylmethyl-8-nitro xanthine;
- 1,3-di-n-butyl-8-amino xanthine;
- 1,3-di-cyclopropylmethyl-8-amino xanthine;
- 1,3-di-cyclobutylmethyl-8-amino xanthine;
- 1,3-di-cyclopentylmethyl-8-amino xanthine;
- 1,3-di-cyclohexylmethyl-8-amino xanthine;
- 1,3-di-cyclopropyl-8-amino xanthine;
- 1,3-di-n-butyl-8-acetamido xanthine;
- 1,3-di-n-butyl-8-chloro xanthine;
- 1,3-di-n-butyl-8-bromo xanthine;
- 1,3-di-cyclopropylmethyl-8-chloro xanthine;
- 1,3-di-cyclohexyl-8-chloro xanthine;
- 1,3-di-n-butyl-8-piperidino xanthine;
- 1,3-di-cyclopropylmethyl-8-morpholino xanthine;
- 1,3-di-n-butyl-8-pyrrolidinyl xanthine;
- 1,3-di-cyclopropylmethyl-8-pyrrolidinyl xanthine;
- 1,3-di-cyclopropylmethyl-8-piperidinyl xanthine;
- 1,3-di-cyclohexylmethyl-8-piperidinyl xanthine;
- 1,3-di-cyclohexylmethyl-8-bromo xanthine; and
- 1,3-di-cyclohexyl-8-nitro xanthine; or if appropriate a pharmaceutically acceptable salt thereof.

16. A process according to claim 2 for preparing 1,3-di-cyclopropylmethyl-8-amino xanthine or a pharmaceutically acceptable salt thereof.

17. A process according to claim 1, which process comprises reacting a compound of formula (II) with a nitrating agent, to provide a compound of formula (IA) wherein R^{3a} represents a nitro group, and thereafter if required carrying out the following optional steps:

- (i) converting any group R^{1a} to R^1 and/or R^{2a} to R^2 ;
- (ii) converting the nitro group into another group R^{3a} ;
- (iii) converting a compound of formula (IA) into a pharmaceutically acceptable salt.

18. A process according to claim 1 wherein the conversion of the nitro group into another group R^{3a} is selected from the list consisting of:

- (i) converting the nitro group into a halogen atom;
- (ii) converting the nitro group into an amine group;
- (iii) converting the nitro group into a halogen atom followed by conversion of the halogen atom into a group $-NR^{4b}R^{5b}$ wherein R^{4b} and R^{5b} together with the nitrogen atom to which they attached form an optionally substituted, heterocyclic group; and
- (iv) converting the nitro group into an amino group and thereafter alkylating and/or acylating the amino group to provide a group $-NR^{4c}R^{5c}$ wherein R^{4c} represents hydrogen, alkyl or alkylcarbonyl and R^{5c} represents alkyl or alkylcarbonyl.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 389 282 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90303093.0

(51) Int. Cl.⁵: C07D 473/06, A61K 31/52

(22) Date of filing: 22.03.90

(30) Priority: 23.03.89 GB 8906792

(43) Date of publication of application:
26.09.90 Bulletin 90/39(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE(86) Date of deferred publication of the search report:
01.07.92 Bulletin 92/27

(71) Applicant: **BEECHAM - WUELFING GmbH & Co. KG**
Stresemannallee 6 P.O. Box 25
W-4040 Neuss(DE)
Applicant: **Beecham Group p.l.c.**
SB House Great West Road
Brentford Middlesex TW8 9BD(GB)


(72) Inventor: **Maschler, Harald, c/o Beecham-Wuelfing GmbH & Co KG, Bethelner Landstrasse W-3212 Gonau (Leine)(DE)**

Inventor: **Spicer, Barbara Ann, Beecham Pharmaceuticals**
Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT 18 5XQ(GB)
Inventor: **Smith, Harry, Beecham Pharmaceuticals**
Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT 18 5XQ(GB)

(74) Representative: **Rutter, Keith et al**
Smith Kline Beecham, Corporate Patents,
Great Burgh, Yew Tree Bottom Road
Epsom Surrey KT18 5XQ(GB)

(54) **Xanthinederivatives, process for their preparation and their pharmaceutical use.**

(57) A method for the treatment of cerebrovascular disorders and/or disorders associated with cerebral senility and/or other disorders which method comprises the administration of an effective, non-toxic amount of a compound of formula (I):



(I)

or if appropriate a pharmaceutically acceptable salt thereof, wherein R¹ and R² each independently represent alkyl or a moiety of formula (a):

-(CH₂)_m-A (a)

wherein m represents zero or an integer 1, 2 or 3;

A represents a substituted or unsubstituted cyclic hydrocarbon radical; and

R³ represents a halogen atom, a nitro group, or a group -NR⁴R⁵ wherein R⁴ and R⁵ each independently represents hydrogen, alkyl or alkylcarbonyl or R⁴ and R⁵ together with the nitrogen to which they are attached forming an optionally substituted, heterocyclic group; certain novel compounds falling within formula (I) and

compositions comprising such compounds.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 30 3093

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
X	BIOCHEMISTRY, vol. 16, no. 15, 26th July 1977, pages 3316-3321, American Chemical Society, Washington, D.C., US; G. L. KRAMER et al.: "Selective inhibition of cyclic nucleotide phosphodiesterases by analogues of 1-methyl-3-isobutylxanthine" * Page 3317, column 2, compound 17 *	2-17	C 07 D 473/06 A 61 K 31/52
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 24, no. 8, August 1981, pages 954-958, American Chemical Society, Washington, D.C., US; J. N. WELLS et al.: "Inhibition of separated forms of cyclic nucleotide phosphodiesterase from pig coronary arteries by 1,3-disubstituted and 1,3,8-trisubstituted xanthines" * Page 956, compounds 5,6; page 958, compounds 5,6 *	2-17	
X	WO-A-8 606 724 (SCHERING CORP.) * Page 2, lines 1-6; page 7, formula II; page 14, sub G *	3-17	TECHNICAL FIELDS SEARCHED (Int. Cl. 5) C 07 D 473/00 A 61 K
X	EP-A-0 152 944 (BOEHRINGER MANNHEIM) * Page 31, lines 26-31; page 32, lines 1-34; claims *	1-27	
A	EP-A-0 267 676 (BEECHAM) * Whole document *	1	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 27-03-1992	Examiner LUYTEN H.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPF FORM 150 01A2 (1/1981)



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet -B-

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

EP 90 30 3093 -B-

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions,

namely:

1. Claims 1(partially), and 2-18 : Purines derivatives, processes for their preparation and pharmaceutical compositions.
2. Claim 1(partially) : Second medical use of known purine derivatives.

